

ESSEX INSTITUTE  
MAY 26 1907  
SALEM, MASS.

VOL. V.




MAY, 1907

NO. 77

# MYCOLOGICAL BULLETIN

W. A. KELLERMAN, Ph. D.

OHIO STATE UNIVERSITY



ENTERED AS SECOND CLASS MATTER MAY 11, 1906, AT THE POSTOFFICE AT COLUMBUS, OHIO.

Edited and Published by  
W. A. KELLERMAN

Press of VANCE-POLAND CO.  
Columbus, Ohio

589.22

M99

CARD  
CATALOGUED.

# Supplies for the Botanist



\$ .30

## Tripod Dissecting Microscope

A convenient and portable instrument for botanists, as it can be used for dissection when mounted, or the lens may be unscrewed and used as a pocket glass. It gives a large, clear image.


## Triple Aplanats

This magnifier consists of a crown lens cemented between



\$3.25

two flint lenses and is thoroughly achromatic. It gives clear, flat images, free from distortion. The mountings are German silver, nickel plated.



DIAMETER, 3-4 IN. \$1.50

## Coddington Magnifier

This magnifier gives clear and perfect definition. The mountings are neatly finished and nickeled.

SEND FOR BOOKLET OF MAGNIFIERS AND READERS

# Bausch & Lomb Optical Co.

Rochester, N. Y.

New York - Boston - Washington - Chicago - San Francisco

When answering advertisement mention Mycological Bulletin.



# You Can Use Engravings

in *your* business, just the same as other men use them. Time was when engravings were really too expensive for common use, but they are cheap now———o

## BUCHER ENGRAVING COMPANY

COLUMBUS,  
OHIO

Portraits, Buildings  
Machinery, Live Stock  
Advertising Cuts  
Stationery Headings  
Cover Pages, Etc.

We have a beautiful specimen book which shows the quality of our work, and we send it free on request

WRITE US ABOUT ANY KIND OF ENGRAVED DESIGN FOR ANY PURPOSE


*We make the engravings for this Publication*

## Standard Botanical Text Books

ANDREWS'S (E. F.) Botany All the Year Round.....	\$1 00
The same. With Brief Flora of the Eastern United States....	1 50
APGAR'S (A. C.) Trees of the Northern United States.....	1 00
APGAR'S (E. A. & A. C.) New Plant Analysis.....	55
CHAPMAN'S Flora of the Southern United States Third Edition....	4 00
COULTER'S Manual of the Botany of the Rocky Mountains.....	1 62
GRAY'S Lessons in Botany .....	94
Outlines of Botany (Leavitt) .....	1 00
The same. With Field, Forest and Garden Flora.....	1 80
The same. With Manual of Botany.....	2 25
Field, Forest and Garden Botany, Flora only .....	1 44
School and Field Book of Botany. Lessons and Flora.....	1 80
Manual of Botany. Flora only.....	1 62
The same. Tourist's Edition.....	2 00
Lessons and Manual. One volume.....	2 16
GRAY'S New Botanical Text-Book:	
Vol. I., Structural Botany (Gray).....	2 00
Vol. II., Physiological Botany (Goodale).....	2 00
GRAY & COULTER'S Text-Book of Western Botany.....	2 16
LEAVITT'S Outlines of Botany .....	1 00
The same. With Gray's Field, Forest and Garden Flora .....	1 80
The same. With Gray's Manual of Botany.....	2 25
WOOD'S (Alphonso) Object Lessons in Botany.....	1 00
Lessons in Botany .....	90
New American Botanist and Florist—Lessons and Flora (Willis) ..	1 75

**AMERICAN BOOK COMPANY, Cincinnati**

When answering advertisement mention Mycological Bulletin.



Nine Mushrooms. Weight four pounds.  
Grown from Pure Culture Spawn.

## Mushrooms.

"Wait till you see it is a good thing, but don't wait till everyone sees it. You will then be too late."

Did you ever think just what is expressed in those words?

Did you ever think of them in connection with mushrooms?

Think of MUSHROOM GROWING—a coming industry in America.

We tell you more about it in our booklet. We also give you some interesting facts about PURE CULTURE SPAWN. They're yours on request.

**PURE CULTURE SPAWN CO.**

609 W. Fourth St.,

Factory: Pacific, Mo.

CINCINNATI, OHIO

---

When answering advertisement mention Mycological Bulletin.



# Mycological Bulletin

No. 77

W. A. Kellerman, Ph. D., Ohio State University

Columbus, Ohio, May, 1907.

---

## THE EDITOR NEEDS LITTLE SPACE.

We print the concluding portion of Mr. Kauffman's admirable Key to the species of *Cortinarius* in this number. It can not fail to be very useful to those who have the courage to attack this difficult genus. Our species are so numerous that every one can take a turn at the Key—but we will not be surprised if some of the amateurs or even students later report to us that some things are easier to master than this Key or the species of *Cortinarius*.

We are able to furnish an additional diversion also in the presentation of matter that amateurs may not have thought of as in the nature of "Mushrooms"—referring to the article on Moulds by Superintendent Sumstine. But these are Fungi, and the word Mycology includes this interesting group. We will be glad for additional articles and notes by patrons of the BULLETIN.

---

## KEY TO THE SPECIES OF *CORTINARIUS*.

(Continued from p. 315.)

- f. Stem marginate-bulbous; gills very narrow and crowded; whole plant violaceous, large ..... *C. Michiganensis* sp. nov.
- ff. Stem subequal or clavate; gills subdistant, adnate; whole plant violaceous-purple, medium size ..... *C. iodes* B. & C.
- ee. Pileus not glutinous
  - f. Flesh and gills turning purple when bruised... *C. purpurascens* Fr.
  - ff. Flesh not turning purple
    - g. Stem marginate-bulbous; pileus yellowish or brownish, tinged violaceous; medium size ..... *C. coerulescens* Fr.

- gg. Stem not marginate-bulbous
  - h. Pileus yellow; gills violaceous to cinnamon; stem white with violaceous apex ..... *C. Berlesianus* Sacc. & Cub. (Syn.=*C. tricolor* Pk.)
  - hh. Pileus and gills lilac; plant small..... *C. croceo-coerulius* (Pers.) Fr.
  - ddd. Pileus with neither olivaceous nor violaceous tints (except the first)
- e. Pileus glutinous
  - f. Gills olivaceous; pileus brownish-ochraceous..... *C. glutinosus* Pk.
  - ff. Gills whitish at first
    - g. Pileus bay-red ..... *C. maculipes* Pk.
    - gg. Pileus pale ochraceous, spores globose... *C. sphaerosporus* Pk.
    - fff. Gills violaceous at first, spores as in preceding... *C. delibutus* Fr.
- ee. Pileus not glutinous
  - f. Stem marginate-bulbous
    - g. Gills at first whitish ..... *C. multiformis* Fr.
    - gg. Gills at first blue ..... *C. glaucopus* Fr.
    - ggg. Gills at first yellow..... *C. fulgens* (Alb. & Schw.)
  - ff. Stem not marginate-bulbous, clavate to subequal
    - g. Gills and stem pallid at first, soon tinged brown
      - h. Pileus watery-cinnamon to brick-red on disk; in woods *C. glabrellus* sp. nov.
      - hh. Pileus whitish to pale clay-color; in mushroom and flower-beds ..... *C. intrusus* Pk.
    - gg. Gills and apex of stem violaceous at first, soon brownish *C. lanatipes* Pk.
    - ggg. Gills and pileus drab-gray; viscid universal veil present *C. sterilis* Kauff.
- B. Cuticle of pileus not composed of gelatinous cells, hence never viscid nor gelatinous. [*Inoloma*, *Talamonia*, *Dermocybe*, and *Hydrocybe*.]
  - a. Spores 12-16 $\mu$  long
    - b. Pileus rather large, squamulose; whole plant dark violaceous *C. violaceus* Fr.
    - bb. Pileus small, chestnut color; stem white; spores 16x11 $\mu$ .... *C. sericipes* Pk.
  - aa. Spores 10-12 $\mu$  long
    - b. Plants small, 2-4 cm. tall
      - c. Pileus hygrophanous, glabrous, bay-red (moist); gills subochraceous ..... *C. badius* Pk.
      - cc. Pileus not hygrophanous, densely fibrillose; gills yellow *C. aureifolius* Pk.
    - bb. Plants larger
      - c. Stem distinctly sheathed or ringed by the universal veil
        - d. Pileus tawny; stem with cinnabar-colored, persistent, concentric rings ..... *C. armillatus* (Alb. & Schw.)
        - dd. Pileus purplish-brown, copper-brown, etc., to drab; stem peronate, i. e., sheathed with a universal veil... *C. torvus* Fr. *C. torvus nobilis* Pk.

- d. Pileus tinged yellow or rufous; stem peronate and annulate by a white universal veil .....*C. canescens* Ph.
- cc. Stem not sheathed or ringed; the universal veil evanescent or absent.
- d. *Pileus hygrophanous, fibrillose-squamulose (like C. paleaceus)*
- e. Pileus dingy chestnut (moist); stem long and slender...*C. gracilis* Ph.
- ee. Pileus grayish; stem stout and short, bulbous.....*C. grieseus* Ph.
- dd. *Pileus not hygrophanous, merely silky or innately fibrillose*
- e. Pileus reddish-gray, tinged purplish; gills purple or violaceous; spores 10-12μ long .....*C. pulchrifolius* Ph.  
   *C. rubrocineruus* Ph.
- ee. Pileus, stem and gills lilac; spores 9-10μ.....*C. lilacinus* Ph.
- eee. Pileus, stem and gills violaceous at first; spores 10-12μ long....  
   *C. rimosus* Ph.
- aaa. Spores 4-9μ long; if longer, plants are whitish or violaceous
- b. Stem and pileus scaly or shreddy
- c. Scales red (scarlet to vermilion).....*C. bolaris* Fr.
- cc. Scales brown to blackish
- d. Plant large, watery-spongy, soon dark chocolate colored.  
   *C. squamulosus* Ph.
- dd. Plants of medium size, wood-brown....*C. pholideus* Fr.
- bb. Stem not scaly.
- c. Stem with more or less persistent annular rings, or peronate
- d. *Plants large, 2-8 cm. or more tall; pileus in proportion*
- e. Pileus watery-cinnamon (moist); gills very distant...*C. distans* Ph.
- ee. Pileus buff, ochraceous, clay-colored or tawny
- f. Gills at first yellow or yellowish
- g. Pileus at first buff; stem peronate by the thin universal veil..  
   *C. flavifolius* Ph.
- gg. Pileus ochraceous to ferruginous; subannulate.....  
   *C. Morrisii* Ph.
- ggg. Pileus at first tawny-yellow, with pointed squamules on disk; peronate by tawny-yellow universal veil.....  
   *C. annulatus* Ph.
- ff. Gills at first brownish or ochraceous; pileus rufous-ochraceous
- g. Spores elliptical.....*C. bivexus* Fr.
- gg. Spores spherical, minute, 4-5μ diameter .....  
   *C. subbivexus* sp. nov.
- eee. Entire plant saffron-yellow.....*C. croceocolor* Kauff.
- eeee. Pileus some shade of blue or purple when young, buff to tan when old
- f. Plants stout, amber-purple to buff; pileus punctate; in or near swamps, in large troops.....*C. umidicola* Kauff.
- ff. Mature plants rather slender; pileus fawn-colored, tinged lavender when young, not punctate; common in hemlock woods....  
   *C. deceptivus* Kauff.
- dd. *Plants small, subannulate; pileus less than 3-4 cm. broad*
- e. Pileus fuscous, covered with white villose fibrils.....  
   *C. paleaceus* (Weimm.) Fr.
- ee. Pileus not villose-squamulose, cinnamon to chestnut color



- f. Gills and stem violaceous at first.....*C. subflexipes* Pk.
- ff. Gills and stem pallid to brownish
  - g. On rotten wood; pileus watery cinnamon..*C. ligniarius* Pk.
  - gg. On ground or moss; pileus bay to chestnut brown; annulus often distinct.....*C. castaneoides* Pk.
  - cc. Stem with no annulus, or annulus evanescent
    - d. *Stem bulbous or clavate*
- e. Bulb depressed-marginate; gills heliotrope purple when young.....*C. obliquus* Pk.
- ee. Bulb clavate to subclavate
  - f. Color of plant lilac to violaceous-white
    - g. Plants of medium size, violet tinge evanescent, never yellowish .....*C. alboviolaceus* (Pers.) Fr.
    - gg. Plants medium to large, lilac tinge persistent..*C. lilacinus* Pk.
    - ggg. Plants medium to small, violaceous to cinereous, tinged yellow or brown.....*C. simulans* Pk.
  - ff. Color of plant deep chrome, unchanging.....*C. callisteus* Fr.
  - fff. Color of plant watery-cinnamon or rufous-cinnamon (moist)
    - g. Stem whitish, pileus rufous-cinnamon to tan; not hygrophanous.....*C. subsalmoneus* sp. nov.
    - gg. Stem red; pileus hygrophanous, pinkish-ochraceous (dry)..*C. rubipes* Kauff.
    - dd. *Stem subequal or tapering downward*
- e. Pileus distinctly hygrophanous
  - f. Plant small; pileus 2 cm. broad or less
    - g. Gills and stem violaceous when young
      - h. Stem stout, smooth; spores 7-9 $\mu$  long.....*C. castaneus* (Bull.) Fr.
      - hh. Stem slender; spores 6-7 $\mu$  long
        - i. Gills and stem pale reddish violaceous at first; pileus blackish-brown; in woods.....*C. subflexipes* Pk.
        - ii. Gills dark-violaceous at first; pileus fuscous, tinged violaceous; on sphagnum.....*C. fuscoviolaceus* Pk.
    - gg. Gills ochraceous, pale; stem whitish, not slender.....*C. pulcher* Pk.
  - ff. Pileus broader than 2 cm.
    - g. Pileus tawny orange to cinnamon; stem pale.....*C. armeniacus* (Schaeff.) Fr.
    - gg. Pileus watery-cinnamon; gills very distant...*C. distans* Pk.
    - ggg. Pileus and stem pale lavender; stem long and attenuated .....*C. everneus* Fr.
- ee. Pileus not hygrophanous
  - f. Pileus chestnut or cinnamon color
    - g. Stem whitish, soon dingy to brownish....*C. castanellus* Pk.
    - gg. Stem yellow, no olivaceous tinge
      - h. Gills at first yellow.....*C. cinnamomeus* (L.) Fr.
      - hh. Gills at first flame scarlet.....*C. semisanguineus flamineus* Kauff.
      - hhh. Gills at first dark blood-red....*C. semisanguineus* Fr.



- ff. Pileus tawny-olive; stem yellow, tinged olivaceous.....  
*C. croceus* Fr.
- fff. Pileus and stem scarlet or blood red
- g. Pileus broad as compared with the rather short stem; spores  
 $8 \times 5\mu$ .....*C. cinnabarinus* Fr.
- gg. Pileus narrow; stem longer; spores  $6 \times 4\mu$ .....  
*C. sanguineus* (Wulf.) Fr.



FIG. 245.—EARTH-STAR. GE-AS'-TER MIN'-I-MA.—A rather common plant here shown in natural size. The cut was made from specimens sent by Supt. M. E. Hard, who collected them near Chillicothe, Ohio, November 9, 1905.

CORRECTION.—The above half-tone was issued on p. 201 as *Geaster triplex*, which was a mistake; it should have been *Geaster minima* as here given. Both of these species were received at the same time and inadvertently the name of the little species got wrong in print.

## MOULDS.

BY DAVID R. SUMSTINE.

It is not necessary to go to the fields and the forest in search of plants. The kitchen, the cupboard, the cellar, the manure heap about the barn, the decaying vegetable matter about the garden or about the house have a distinctive flora as interesting as the flora of field and forest. The plants of the latter flora are more conspicuous, but the plants of the former flora are just as beautiful and have just as interesting life history and in the economy of nature are just as useful.


There are many genera and species represented in this flora but only a few belonging to the family *Mucoraceae* will be discussed at this time.

These plants are commonly called *Moulds*, but all the so-called Moulds do not properly belong to this family. This is especially true of the common green mould found on canned fruit. It belongs to an entirely different family.

A piece of bread or sweet potato laid in a moist place for a few days will produce plenty of specimens for study. It seems that the spores of moulds float in the air and fall upon various substances. Whenever the proper conditions of temperature and moisture are supplied the spores germinate and develop. The spores retain the power of germination for a long time.


A little observation will show the thread like mycelium spreading in and upon the substratum. At different parts of the mycelium *sporangio-phores* arise. At the top of the sporangiophores are developed the sporangia containing spores.

The following species can be found almost anywhere during the summer months:



*Mucor mucedo*. a Sporangiophores. b Spores.





a Spinulose mycelium of *Mucor fusiger*. b Spores of same. c Columella of *Mucor stercoreus*. d Spores of same. e Columella of *Phycomyces nitens*. f Spores of same.

FIG. 247. ILLUSTRATIONS OF MOULDS.

### MUCOR.

This genus is characterized by simple or branched, but not dichotomously branched, sporangiophores. The membrane of the sporangium is not cuticularized and soon disappears. The sporangia are of one kind and are furnished with columella.

#### MUCOR MUCEDO LINNAEUS.


The mycelium is white at first, then brown or even black. The sporangiophores are usually in clusters of three or more, reaching 5 mm. in height. The columella is very prominent, but finally collapses. The spores are globose, oval or irregular. This is the common black mould of bread. The following names are also given to this species, *Mucor stolonifer*, *Rhizopus nigricans*.

#### MUCOR STERCOREUS (TODE) LINK.

The sporangiophores are erect, undivided, gray; the sporangia are large, yellowish, brownish when old; the membrane disappears and leaves a collar at the base; the columella is variously shaped, cylindrical or conical; spores somewhat elliptical. It grows on dung. *Mucor mucedo* Fresenius has also been applied to this mould.

#### MUCOR FUSICER LINK.

The sporangiophores are simple, erect, bulbous below, but attenuated upwards, gray to brown in color; columella is large; spores spindle shaped; the aerial mycelium is divided into thorny branches. It grows on Agarics.



a Sporangiophore of *Pilobolus crystallinus*. b Spores of same.  
c Sporangiophores of *Sporodinia aspergillus*. d Spores of same.

FIG. 248. ILLUSTRATIONS OF MOULDS.

#### PHYCOMYCES.

In general appearance the species of this genus resemble the species of *Mucor*. The chief difference is in the formation of zygospores. The metallic appearance will generally separate it from *Mucor*.

*PHYCOMYCES NITENS* (AGARDH) KUNZE.

The sporangiophores are simple, olive brown, very large, 7-30 cm. long; the sporangia are round, large; columella, pear shaped; spores ellipsoid. It grows on oily substances. Ground flax seed makes a good medium for cultivation.

#### PILOBOLUS.

This genus is easily recognized by the peculiar formation of the sporangium whose upper part is cuticularized. The sporangiophore is very much enlarged or distended right below the sporangium.

*PILOBOLUS CRYSTALLINUS* (WIGGERS) TODE.

This is common on horse dung. The sporangium is black and at maturity is thrown off with considerable force; the spores are colorless or yellowish.

#### SPORODINIA.

This is a monotypic genus and is known by the dichotomously branched sporangiophores.

*SPORODINIA ASPERGILLUS* (SCOPOLI) SCHROETER.


The sporangia are variously colored, but usually brown or black at maturity. The spores are round or ellipsoid. It grows on Boleti and Agarics.

All figures were drawn by Stella Sumstine, not to any definite scale.



# NEW SPENCER MICROSCOPE

No. 404



No. 40 Stand  
Abbe Condenser  
Two Iris Diaphragms  
Triple Nosepiece  
16 mm. (1-3 ") Objective  
4 mm. (1-6 ") Objective  
2 mm. (1-12") Oil Im-  
mersion Objective  
Two Eyepieces  
Mahogany Cabinet

**THE BEST**  
**MICROSCOPE EVER OFFERED FOR**  
**\$75.00 BECAUSE:**

The lenses are unsurpassed

The Stand is the product of the best of material and workmanship

The coarse and fine adjustments are the best made

The fine adjustment is protected by a neat handle provided for carrying the instrument

The stage is large—67 mm. free distance from the optical axis to the base of the arm is the greatest yet produced

The stage is completely covered with vulcanite from top to bottom

**SPENCER LENS COMPANY**  
**BUFFALO, N. Y.**

When answering advertisement mention Mycological Bulletin.

## FOR SALE

Half-tone Portraits of the following eminent Mycologists:	
Peck, Ellis, Arthur, Burrill, Farlow, Tracy, Earle, Atkinson, Holway, Thaxter, Clinton, Magnus, Oudemans, Patouillard, Hennings, Rehm, Cooke, Saccardo, Dietel, Sydow, P., Sydow, H., Hollis, Bresadola, Massee, Bubak, Salmon, each .....	10 cts.
Mycological Bulletin, Vol. I, 1903.....	50 cts.
“ “ “ II, 1904.....	50 cts.
“ “ “ III, 1905.....	50 cts.
“ “ “ IV, 1906.....	50 cts.
“ “ Current volume.....	25 cts.
Mycological Glossary .....	25 cts.
North American Species of Marasmius (a Monograph by Morgan).....	50 cts.
North American Species of Lepiota (a Monograph by Morgan).....	50 cts.
Journal of Mycology, Vol 8—1902.....	\$2 00
“ “ “ 9—1903.....	2 00
“ “ “ 10—1904.....	2 00
“ “ “ 11—1905.....	2 00
“ “ “ 12—1906.....	2 00
“ “ “ Current Vol.....	2 00

W. A. Kellerman, Columbus, Ohio

**2nd Edition** of the book which is “without doubt the most important and valuable work of its kind.”—*Plant World*.

# MUSHROOMS

By Prof. GEORGE FRANCIS ATKINSON, of Cornell

Edible, Poisonous Mushrooms, Etc. With recipes for cooking by Mrs. S. T. Rorer, and the chemistry and toxicology of Mushrooms, by J. F. Clark. With 230 illustrations from photographs, including fifteen colored plates by F. R. Rathbun, 320 p. 8vo, \$3.00 net (by mail \$3.23.)

Among the additions in this second edition of Prof. Atkinson's remarkable book are ten new plates, chapters on “The Uses of Mushrooms” and on “The Cultivation of Mushrooms” illustrated by several flashlight photographs.

THE NATION said of the first edition “If Prof. Atkinson would fill in the gaps, and 20 or 30 plates and issue a second edition . . . he would win the gratitude of every amateur and professional mycologist in the country. (This second edition is the one now issued.)

Prospectus of New American Nature Series Free on Application.

**HENRY HOLT & COMPANY 29 W. 23rd St., NEW YORK**

When answering advertisement mention Mycological Bulletin.



LAMBERT'S PURE CULTURE

# Mushroom

---

---

# Spawn

---

---



Produced by the new grafting process from selected and prolific specimens, thoroughly acclimatized, has *never failed to run*.

This spawn is made from cultures taken by the *selective* method, recently discovered, from choice specimens of the best varieties of mushrooms known to be thoroughly acclimatized, and selected with special reference to their size, flavor, vigor and prolificness. The elements of uncertainty surrounding the old *chance* method of producing wild spawn (English and French), is therefore eliminated, and a uniform crop of the best marketable variety is thus assured

to the exclusion of all other and inferior fungi.

The importance of this discovery will be realized from the mere statement that, for the first time in the history of mushroom culture, can spawn be offered of a specific variety of mushrooms.

Of the varieties already developed, we are offering pure culture spawn of the *Agaricus Campestris* (white, cream or brown), *A. Arvensis* (cream), and *A. Villaticus* (cream).

FOR SALE BY ALL LEADING SEEDSMEN.

Practical Instructions on "MUSHROOM CULTURE"  
mailed free on application.

## American Spawn Company,

ST. PAUL, MINN.

When answering advertisement mention Mycological Bulletin.



This space is owned by  
THE TERRY ENGRAVING CO.  
COLUMBUS, OHIO



When answering advertisement mention Mycological Bulletin.